



CHECK YOUR ANSWERS

- (c) Find the time taken for the car to travel from point A to point C . (2)

This image shows a full page of white paper with horizontal blue ruling lines. The lines are evenly spaced and run across the width of the page. In the bottom-left corner, there is a small, stylized blue gear-like icon.

2

In the model of the motion, the cyclist starts from rest at point C when $t = 0$, then moves with a constant acceleration of 1.2 ms^{-2} . When $t = 9$, the cyclist arrives at point D .

- (c) Find the time taken to reach the midpoint of C and D . (3)

(Total for Question 2 is 7 marks)

3

When $t = 24$, the velocity of the train is 34 ms^{-1}

- (c) Find the distance travelled by the train in the 30 second journey. (3)

(Total for Question 3 is 7 marks)



4 A cyclist travels along a straight horizontal road.

In the model of the motion, the cyclist is travelling at 9 ms^{-1} when they apply the brakes, causing them to decelerate uniformly. The cyclist comes to rest after travelling 27 m with the brakes applied.

- (b) Find the time taken to come to rest whilst they are braking. (2)

[illegible]

(Total for Question 4 is 4 marks)

5 A bus travels along a straight horizontal road.

In the model of the motion, the bus accelerates uniformly from a velocity of 8.5 ms^{-1} to a velocity of 16.5 ms^{-1} , in 5 seconds. The bus then maintains a constant velocity of 16.5 ms^{-1} for a further 15 seconds.

- (b) Find the total distance travelled by the bus during the 20 second period. (4)



(Total for Question 5 is 6 marks)



6

In the model of the motion, the runner starts from rest when $t = 0$ before travelling with constant acceleration 1.6 ms^{-2} , until they reach a velocity of 6.4 ms^{-1}

The runner then maintains a constant velocity of 6.4 ms^{-1} until they have travelled a total distance of 500 m.

Show that it takes the runner more than 80 seconds travel the distance of 500 m.

(Total for Question 6 is 4 marks)

7

In the model of the motion, the motorcycle travels with constant acceleration between points A and B , passing point A with velocity 18 ms^{-1} and point B with velocity 32 ms^{-1}

Given that the distance between A and B is 125 m, find

(a) the acceleration of the motorcycle (2)

(b) the time taken for the motorcycle to travel between points A and B . (2)

(Total for Question 7 is 4 marks)



9 An object travels in a straight line with constant acceleration $a \text{ ms}^{-2}$ for t seconds.

It has initial velocity $u \text{ ms}^{-1}$ and final velocity $v \text{ ms}^{-1}$

Given that $v = u + at$ and $s = \frac{1}{2}(u + v)t$ show that

$$(a) \quad s = ut + \frac{1}{2}at^2 \quad (3)$$

$$(b) \quad s = vt - \frac{1}{2}at^2 \quad (3)$$

$$(c) \quad v^2 = u^2 + 2as \tag{3}$$



6

(Total for Question 9 is 9 marks)

10 A train travels along a straight horizontal track.

In the model of the motion, when $t = 0$ the train has travels with uniform acceleration 0.3 ms^{-2} for T seconds, from an initial velocity of 15 ms^{-1} to a velocity of $V \text{ ms}^{-1}$.

At time $t = T$ the train has travelled a total distance of 255 m.

Find the values of V and T .

(Total for Question 10 is 4 marks)

11 A car travels along a straight horizontal road.

In the model of the motion,

- the car travels with constant acceleration
- when $t = 2$, the car has velocity 10 ms^{-1}
- when $t = 8$, the car has travelled a total distance of 104 m

Find the velocity of the car when $t = 9$

(Total for Question 11 is 6 marks)



12 A cyclist travels along a straight horizontal road between points A and B .

In the model of the motion, at $t = 0$ the cyclist passes point A , travelling at 12 ms^{-1} and then accelerates uniformly at 0.5 ms^{-2} for T seconds, reaching a velocity of $V \text{ ms}^{-1}$ when they pass point B .

The distance between A and B is D metres

In the final 2 seconds of the journey the cyclist travels a distance of 34 metres.

Find the values of V , T and D .

[illegible]

(Total for Question 12 is 6 marks)

